

## STA 3000, Fall 2009, Assignment #1, Answer to Question 1

**Question 1:** For the model  $Y \sim P_\theta$ , suppose that  $s(Y)$  is a sufficient statistic. Prove that for the model  $Y_1, \dots, Y_n \stackrel{\text{iid}}{\sim} P_\theta$ , the order statistics,  $S_{(1)}, \dots, S_{(n)}$ , of  $S_1, \dots, S_n$ , where  $S_i = s(Y_i)$ , are sufficient. Prove this directly from the (classical) definition of a sufficient statistic, without using any theorems about sufficient statistics.

**Answer:** We need to prove that the conditional distribution of  $Y_* = (Y_1, \dots, Y_n)$  given  $S_{(*)} = (S_{(1)}, \dots, S_{(n)})$  is the same for all  $\theta$ . I will assume that we can naively express conditional distributions by conditional densities that are ratios of joint densities. This will certainly be fine when the  $Y_i$  are discrete. The density function for the original model for a single  $Y$  will be written as  $f_Y$ , and the density function for  $s(Y)$  will be written as  $f_S$ .

To start,

$$\begin{aligned} f_{Y_*|S_{(*)}}(y_*|s_{(*)}) &= \frac{f_{Y_*, S_{(*)}}(y_*, s_{(*)})}{f_{S_{(*)}}(s_{(*)})} \\ &= \frac{f_{Y_*}(y_*) I(\text{there is a permutation } \pi \text{ such that for all } i, s_{(i)} = s(y_{\pi(i)}))}{f_{S_{(*)}}(s_{(*)})} \end{aligned}$$

We can now use independence of the  $Y_i$  to write  $f_{Y_*}(y_*) = \prod_i f_Y(y_i)$ . Using independence of the  $S_i$ , we can write

$$f_{S_{(*)}}(s_{(*)}) = \sum_{\substack{s_1, \dots, s_n \\ \text{permuting} \\ s_{(1)}, \dots, s_{(n)}}} \prod_i f_S(s_i) = N(s_{(*)}) \prod_i f_S(s_{(i)})$$

where the sum is over distinct permuted sequences of  $s_{(1)}, \dots, s_{(n)}$ , and  $N(s_{(*)})$  is the number of such distinct permutations (which will be  $n!$  if the  $s_{(i)}$  are all distinct). With these substitutions, we get

$$\begin{aligned} f_{Y_*|S_{(*)}}(y_*|s_{(*)}) &= \frac{\prod_i f_Y(y_i)}{N(s_{(*)}) \prod_i f_S(s_{(i)})} I(\text{there is a permutation } \pi \text{ such that for all } i, s_{(i)} = s(y_{\pi(i)})) \\ &= \frac{\prod_i f_Y(y_{\pi(i)})}{N(s_{(*)}) \prod_i f_S(s_{(i)})} I(\text{there is a permutation } \pi \text{ such that for all } i, s_{(i)} = s(y_{\pi(i)})) \\ &= \left[ \prod_i \frac{f_Y(y_{\pi(i)})}{f_S(s_{(i)})} \right] \frac{I(\text{there is a permutation } \pi \text{ such that for all } i, s_{(i)} = s(y_{\pi(i)}))}{N(s_{(*)})} \end{aligned}$$

where if the indicator function  $I(\dots)$  above is one, the  $\pi$  appearing in  $y_{\pi(i)}$  is one of the permutations for which  $s_{(i)} = s(y_{\pi(i)})$  for all  $i$  (and otherwise is any arbitrary permutation).

When  $s_{(i)} = s(y_{\pi(i)})$ , we can write  $f_Y(y_{\pi(i)}) = f_{Y,S}(y_{\pi(i)}, s_{(i)})$ , and we can therefore also write  $f_Y(y_{\pi(i)})/f_S(s_{(i)}) = f_{Y|S}(y_{\pi(i)}|s_{(i)})$ , which does not depend on  $\theta$  since  $S$  is sufficient for the model of a single observation,  $Y$ .

The second factor above — the ratio of  $I(\dots)$  and  $N(s_{(*)})$  — does not depend on  $\theta$ . When this factor is not zero, the first factor — the product over  $i$  of  $f_Y(y_{\pi(i)})/f_S(s_{(i)}) = f_{Y|S}(y_{\pi(i)}|s_{(i)})$  — also does not depend on  $\theta$ . It follows that  $f_{Y_*|S_{(*)}}$  does not depend on  $\theta$ , and therefore  $S_{(*)}$  is sufficient.